

## EVALUATION OF THE FUNGITOXIC ACTIVITY OF VEGETAL EXTRACTS ON THE MYCELIAL GROWTH OF PHYTOPATHOGENS

### AVALIAÇÃO DA ATIVIDADE FUNGITÓXICA DE EXTRATOS VEGETAIS SOBRE O CRESCIMENTO MICELIAL DE FITOPATÓGENOS

Talita Pereira de Souza FERREIRA<sup>1</sup>; Gil Rodrigues dos SANTOS<sup>1</sup>;  
Tatiani Pereira de Souza FERREIRA<sup>1</sup>; Tallyta Santos TEIXEIRA<sup>1</sup>; Ryhára Dias BATISTA<sup>1</sup>;  
Raiana Batista da LUZ<sup>1</sup>; Luiz Gustavo de Lima GUIMARÃES<sup>2</sup>

1. Universidade Federal do Tocantins, Gurupi, TO, Brasil. cupufer@gmail.com; 2. Universidade Federal de São João Del Rei, São João Del Rei, MG, Brasil.

**ABSTRACT:** Although the productivity of common bean in Tocantins is economically favorable, it has been infected by various pathogens found in soil. Among the major diseases is the web blight and root rot caused by *Rhizoctonia solani* and collar rot caused by the fungus *Sclerotium rolfsii*. This study aimed to evaluate the fungitoxic activity of methanol extracts of eight plant species on the inhibition of mycelial growth of *S. rolfsii* and *R. solani*. The fungitoxic activities were carried out over the inhibition of mycelial growth by means in vitro assays. The extracts were applied in concentrations of 250, 500, 1000, 2500 e 5000 µg ml<sup>-1</sup> in PDA culture medium. In bioassays, it was found the significant effect of plant, concentration and also their interaction on the antifungal activity of the extracts. However, some extracts showed no inhibition of mycelial growth of the pathogens studied. Among those who had higher inhibitions is the extract of *Lantana trifolia*, which inhibited the mycelial growth of *S. rolfsii* in all concentrations, being the same as 97% for the highest concentration. When the methanol extract of *Piper amplum* Kunth, inhibition of the highest concentration was 83% for *S. rolfsii* and 74% for *R. solani*. These results show the potential of methanolic extract of *Lantana trifolia* and *Piper amplum* Kunth in the control set of plant pathogens studied.

**KEYWORDS:** Methanol extract. *Sclerotium rolfsii*. *Rhizoctonia solani*. *Phaseolus vulgaris*.

#### INTRODUCTION

The common bean (*Phaseolus vulgaris*) is the most important legume crop consumed by humans worldwide. Various pathogens of epidemiological importance can damage common bean crops reducing their quality (PARSA et al., 2016). The risk of losses caused by pests and diseases in bean decreases the profitability for producers, resulting in decreased exploitation of this legume by large producers (GONÇALVES-VIDIGAL et al., 2007; CUNHA et al., 2005). Therefore, there is a growing interest in discovering plant-derived anti-microbial substances as alternative agents against fungal growth (EL-MOUGY et al., 2007).

Of the 60 diseases that can affect bean crops, 31 are caused by fungi. These pathogens are reported as the primary causative agents of diseases in the economically important vegetable crops. Each year they destroy a third of agricultural production worldwide resulting in approximately 20% reduction in yield of cultivars (GARCIA et al., 2002). Among the diseases caused by pathogenic fungi in bean crops, there are the collar rot and the web blight which are caused by the fungi *Sclerotium rolfsii* and *Rhizoctonia solani*, respectively.

So, to avoid significant losses caused by diseased plants, several control methods are currently employed. The most common being the use of synthetic fungicides (BAJWA; KHALID; CHEEMA, 2003). However, the abusive use of these compounds has been causing environmental damage and has even been proven harmful for consumers, besides promoting resistance of the pathogenic microorganisms. Furthermore, these synthetic fungicides also affect the symbiotic microbial population present in the local ecosystem (CUNICO et al., 2004). In view of this, ongoing research is being conducted in order to create and/or discover new methods to halt the advance of pathogenic fungi growth which are environmentally friendly and effective in the action.

Extracts from different parts of medicinal plants have been the source of many compounds that have antimicrobial activities. Many studies have reported the fungitoxic activity of various plant extracts. For instance, Vogt et al. (2012), while studying the effects of the extract of a medicinal plant from Argentina, *Larrea divaricata*, corroborated its empirical use in the treatment of some plant diseases and showed high rate of antifungal activity, making it a potential natural source for biological control. Fiori et al. (2000) reported that the extracts from the leaves of

*Eucalyptus citriodora* inhibited the mycelial growth of the pathogenic fungus *Didymella bryoniae*, while the plant extract of *Achillea millefolium* inhibited the germination of spores of the same fungus. Also in this work, the plant extract *Ageratum conyzoides* both inhibited the growth of mycelial and spore germination of the fungus *Didymella bryoniae*.

Given the above, this work aimed to study the fungitoxic activity of methanol extracts of eight plant species on the inhibition of mycelial growth *Sclerotium rolfsii* and *Rhizoctonia solani*, in order to develop alternative and effective means to control the collar rot, web blight and root rot caused by these pathogens in the culture of the common bean.

## MATERIAL AND METHODS

### Collection and extraction of vegetal plant extracts

Eight plant species were studied: 'sucupira' (*Pterodon emarginatus*), 'mastruz' (*Chenopodium ambrosioides*), mint (*Mentha piperita*), terramycin (*Alternanthera dentata*), 'negramina' (*Siparuna guianensis* Aublet), pepper (*Piper amplum* Kunth), 'camará' (*Lantana trifolia* L.) and 'grandiúva-d'anta' (*Psychotria leiocarpa* Cham. Et Schlechtend). After harvesting, the plant material was sent to the Chemistry Laboratory on the UFT Campus in Gurupi, where they underwent drying and processes to obtain the extracts. The leaves were dried in the shade at room temperature and subsequently were cut and submitted to extraction with cold methanol. Thirty grams of dried leaves were used with 1.5 liters of solvent for each extraction for a period of 7 days. After this period, the mixture was filtered and evaporated under reduced pressure to obtain the extracts (COSTA et al., 2008).

### Isolation of fungi used in biological assay

The pathogens *S. rolfsii* and *R. solani* were isolated from plants of the common bean *Phaseolus vulgaris* with symptoms of the disease (collar rot and root rot) as follows: plant parts with symptoms of the disease were cut and washed in water. Then these pieces were surface sterilized by successive immersion in 70% ethanol for 30 seconds, a solution of 1% sodium hypochlorite for 40 seconds and three times in demineralized water. After being washed several times, they were transferred with sterile tweezers to a Petri plate containing 15 ml of PDA culture medium (Potato Dextrose Agar) and 1 mg of ampicillin. Daily, the Petri dishes were monitored and fungal colonies that showed no contaminants

were transplanted into new plates with the same culture medium (VALADARES et al. 2008).

### Experimental design and biological activity


Isolated pure cultures of plant pathogens *S. rolfsii* and *R. solani* that were 7 days old were separately tested to evaluate fungitoxicity. Bioassays were conducted by adding to the PDA culture medium (Sigma-Aldrich®), aqueous solutions of methanol extracts from the eight lyophilized medicinal plants, examined in appropriate volumes to obtain concentrations of 250, 500, 1000, 2500 and 5000 µg mL<sup>-1</sup>, equally for each microorganism. With the medium already poured into Petri dishes, discs of agar-mycelium of 5 mm diameter made of pure cultures of fungi were inoculated on the surface of each culture with their respective treatments. Subsequently, the plates were sealed with plastic film and incubated in a growth chamber (25 ± 1 °C) and exposed to a photoperiod of 12 hours. To estimate the efficiency of the treatments, the diameter of each of the colonies was measured during the mycelial growth and compared to that of absolute witness. It was observed that it completely covered the surface of the medium. The radial growth was measured with calipers in two orthogonal axes to each other by calculating an average for each plate. The experimental design was completely randomized with three replications. Treatments were arranged in a factorial scheme 2 x 8 x 5 for evaluation as antifungal activity being constituted by combinations of the eight plant species, two pathogenic fungi and concentrations of extracts. Analysis of variance was carried out to check the effects of treatments, using regression analysis to assess the percentage of inhibition of mycelial growth of phytopathogenic fungi, in relation to concentration using the statistical program SISVAR.

## RESULTS AND DISCUSSION

According to the results of the assessment of the fungitoxic activity of plant extracts from eight plant species studied, there was a significant effect on plant factors, concentration, and also the interaction of these factors on antifungal activities presented by the extracts and applied to each test performed. Through this interaction, we observed the existence of a variation of antifungal activity, depending on the extract of each plant and the concentration of each. The values of antifungal activity in the plant extracts in relation to their concentrations are assessed by inhibition assay in

percentages for the two fungi *S. rolfsii* and *R. solani*

are shown in Figure 1.



**Figure 1.** Fungitoxic activity of the methanol extracts of the ten plant species on mycelial growth of pathogenic fungi *Sclerotium rolfsii* and *Rhizoctonia solani*. A: *Piper amplum*; B: *Lantana trifolia*; C: *Mentha piperita*; D: *Chenopodium ambrosioides*; E: *Psychotria leiocarpa*; G: *Siparuna guianensis*; H: *Pterodon emarginatus*; I: *Alternanthera dentata*.

The effect of the extract *L. trifolia* (Figure 1B) against *R. solani* was observed when the witness completely covered the Petri dish. The mycelial growth was reduced at concentrations of 1000, 2500 and 5000  $\mu\text{g ml}^{-1}$ , and there was inhibition at the highest concentration of 64%. Already with the fungus, *S. rolfssii*, the methanol extract showed this inhibitory effect at all concentrations used. An inhibition of 97% was observed in the highest concentration.

Likewise, the extract of *Psychotria leiocarpa* (Figure 1E) significantly reduced the growth of mycelia of both fungi studied. The inhibition was verified from concentration of 5000  $\mu\text{g ml}^{-1}$ , while the highest concentration obtained 72% mycelial inhibition in *S. rolfssii* and 55% in *R. solani*. When these same fungi were exposed to the extract *Piper amplum* (Figure 1A), inhibition of the highest concentration was 83% for *S. rolfssii* and 75% for *R. solani*, considering the inhibitions were observed from lower concentrations.

According to the regression analyses for the two pathogenic fungi, mycelial inhibition in relation to concentrations of methanol extracts of *L. trifolia* (Figure 1B) and of *P. leiocarpa* (Figure 1E), which follow equations of the second order. Already having considered the concentrations of the extract of *P. amplum* (Figure 1A), which follow equations of the first order.

The assessment of the potential fungitoxic plant extract *A. dentata* (Figure 1H), showed no effect on the inhibition of mycelial pathogen *S. rolfssii*. On the other hand, the pathogen *R. solani* was sensitive to all extracts of plants assessed. But some extracts even at the highest concentration showed lower percentage inhibition of mycelial growth even though the extracts; *P. emarginatus* (Figure 1G), *C. ambrosioides* (Figure 1D), *S. guianensis* (Figure 1F) and *A. dentata* (Fig. 1H), exhibited the following activities: 24, 24, 14 and 20%, respectively. Considering the fact that concentration varies according to the regression analyses, the mycelial inhibition of these extracts followed equations of the second degree, except for the inhibitions caused by the different concentrations of the extract *A. dentata* (Figure 1H), which followed an equation of first degree. The extracts of *M. piperita* (Figure 1C), *C. ambrosioides* (Figure 1D), *S. guianensis* (Figure 1F) and *P. emarginatus* (Figure 1G) also had no significant effects on the inhibition of mycelial growth of phytopathogenic fungus *S. rolfssii*, since before their concentration was increased, the mycelial growth of this microorganism was inhibited by 36, 4, 6 and 10%, respectively. However, the methanol extract of

*M. piperita* (Figure 1C) on *R. solani* showed a significant inhibition of 51% at its highest concentration.

Many studies have presented results which corroborate those found in this work, as conducted with plant extracts of *Ocimum basilicum* (basil), *Ruta graveolens* (rue), *Baccharis trimera* (gorse), which showed antifungal activity on *Rhizoctonia solani*, *Sclerotium rolfssii*, *Alternaria alternata*, *Phytophthora* sp. and *Colletotrichum graminicola* (STARGALIN et al., 1999). Significant effects were observed in a concentration of 1% applied to the turmeric extract which has led to a greater than 50% inhibition of growth of *Fusarium oxysporum* (61%) and *R. solani* (61%) (AMARAL; BARA, 2005). Thangavelu et al. (2004) showed that extracts of *Solanum torvum*, *Jatropha glandulifera* and *Embllica officinalis* were highly inhibitory to mycelial growth of *Colletotrichum musae*, such inhibitions being directly related to the concentration factor. Extracts of leaves and seeds of *Pithecellobium dulce* had fungitoxic effects in more than 50% mycelial growth and sporulation of *Botrytis cinerea*, *Penicillium digitatum* and *Rhizopus stolonifer*, the strawberry fruit (BAUTISTA-BAÑOS et al., 2003).

## CONCLUSIONS

Among the methanol extracts of the plants studied, the one showing the best antifungal effect on mycelial growth of the fungus *S. Rolfssii* was the methanol extract of the plant *L. trifolia* with inhibition of 97%.

For *R. Solani*, the methanol extract of *P. amplum* showed the best result among all extracts evaluated, inhibiting 75% growth of mycelia. Furthermore, observing the percentages of inhibition of the two fungi in general, *S. rolfssii* was more resistant to plant methanol extracts than all pathogens studied.

These results show that the potential of methanol extracts for *L. trifolia* and *P. amplum* in the control group of plant pathogens studied. However, it is necessary to conduct further studies employing fractionation on the extract, for identification of molecule(s) performing such activities.

## ACKNOWLEDGEMENTS

This work was supported by the Conselho Nacional de Desenvolvimento Científico e Tecnológico – CNPq – Brazil.

**RESUMO:** Apesar da produtividade do feijão comum no Tocantins ser economicamente favorável, o mesmo pode ser infectado por vários patógenos habitantes do solo, dentre as principais doenças encontra-se a mela e a podridão radicular causadas pelo fungo *Rhizoctonia solani* e a podridão do colo causada pelo fungo *Sclerotium rolfsii*. Este trabalho teve como objetivo avaliar a atividade fungitóxica dos extratos metanólicos de oito espécies vegetais sobre a inibição do crescimento micelial de *Sclerotium rolfsii* e *Rhizoctonia solani*. As atividades fungitóxicas foram realizadas perante a inibição do crescimento micelial por meio de ensaios *in vitro*, sendo os extratos aplicados nas concentrações de 250, 500, 1000, 2500 e 5000 µg ml<sup>-1</sup> em meio de cultura BDA. Observou-se o efeito significativo dos fatores planta, concentração e também da interação destes sobre as atividades fungitóxicas. No entanto, alguns extratos não apresentaram inibição do crescimento micelial dos fitopatógenos estudados. Entre os que apresentaram maiores inibições encontra-se o extrato de *Lantana trifolia*, que inibiu o crescimento micelial do *S. rolfsii* em todas as concentrações, sendo o mesmo de 97% para a maior concentração. Já a concentração mais elevada do extrato metanólico de *Piper amplum* apresentou inibição de 83% sobre o crescimento micelial de *S. rolfsii* e 74% sobre o crescimento micelial de *R. solani*. Tais resultados evidenciam a potencialidade dos extratos metanólicos das folhas de *Lantana trifolia* e de *Piper amplum* no controle dos fitopatógenos estudados.

**PALAVRAS-CHAVE:** Extrato metanólico: *Sclerotium rolfsii*. *Rhizoctonia solani*. Atividade fungitóxica.

## REFERENCES

- AMARAL, M. F. Z. J.; BARA, M. T. F. Avaliação da atividade antifúngica de extratos de plantas sobre o crescimento e fitopatógenos. **Rev. Eletr. de Farm.**, Goiânia, v. 2, (Supl.), n. 2, p. 5- 8, 2005.
- BAJWA, R.; KHALID, A.; CHEEMA, T. Antifungal activity of allelopathic plant extracts III: growth response of some pathogenic fungi to aqueous extract of *Parthenium hysterophorus*. **Pak. J. Plant Pathol.** New York, v. 2, n. 3, p. 145-156, 2003.
- BAUTISTA-BAÑOS, S.; GARCÍA-DOMÍNGUEZ, E.; BARRERA-NECHA, L. L.; REYES-CHILPA, R.; WILSON, C. L. Seasonal evaluation of the postharvest fungicidal activity of powders and extracts of huamuchil (*Pithecellobium dulce*): action against *Botrytis cinerea*, *Penicillium digitatum* and *Rhizopus stolonifer* of strawberry fruit. **Postharvest Biol. Tec.**, Washington, v. 29, p. 81-92, 2003.
- COSTA, F. J.; BANDEIRA, P. N.; ALBURQUERQUE, M. R. J. R.; PESSOA, O. D. L.; SILVEIRA, E. R.; BRAZ-FILHO, R. Constituintes químicos de *Vernonia chalybaea* Mart. **Quim. Nova**, São Paulo, v. 31, n. 7, p. 1691-1695, 2008.
- CUNHA, J. P. A. R.; TEIXEIRA, M. M.; VIEIRA, R. F. Avaliação de pontas de pulverização hidráulicas na aplicação de fungicida em feijoeiro. **Cienc. Rural**, Santa Maria, v. 35, n. 5, p. 1069-1074, 2005. <https://doi.org/10.1590/S0103-84782005000500013>
- CUNICO, M. M.; CARVALHO, J. L. S.; SILVA, V. C.; MONTRUCCHIO, D. P.; KERBER, V. A.; GRIGOLETTI JÚNIOR, A.; AUER, C. G.; MIGUEL, M. D.; MIGUEL, O. G. Avaliação antifúngica de extratos obtidos de *Ottomiamartiana* Miq. (Piperaceae) sobre três fitopatógenos. **Arq. Inst. Biol.**, São Paulo, v. 71, (Supl.), p. 1-749, 2004.
- EL-MOUGY, N. S.; EL-GAMAL, N. G.; ABDEL-KADER, M. M. Control of wilt and root rot incidence in *phaseolus vulgaris* L. by some plant volatile compounds. **J. Plant Prot. Res.**, v. 47, n. 3, p. 255-265, 2007.
- FIORI, A. C. G.; SCHWAN-ESTRADA, K. R. F.; STANGARLIN, J. R.; VIDA, J. B.; CRUZ, M. E. S.; PASCHOLATI, S. F. Antifungal activity of leaf extracts and essential oils of some medicinal plants against *Didymella bryoniae*. **J. Phytopathol.**, Malden, v. 148, p. 483-487, 2000.
- GARCIA, L. C.; JUSTINO, A.; RAMOS, H. H. Análise da pulverização de um fungicida na cultura do feijão em função do tipo de ponta e do volume aplicado. **Bragantia**, Campinas, v. 61, n. 3, p. 291-295, 2002. <https://doi.org/10.1590/S0006-87052002000300011>

GONÇALVES-VIDIGAL, M. C; BONETT, L. P; FILHO, P. S. V; GONELA, A; RIBEIRO, A. S. Genetic control in the performance of common bean differential cultivars to *Colletotrichum lindemuthianum* races. **Arq. Biol. Tecnol.**, Curitiba, v. 50, n. 4, p. 579-586, 2007.

PARSA, S. GARCÍA-LEMO, A. M.; CASTILLO, K.; ORTIZ, V.; LOPEZ-LAVALLE, L. A. B.; BRAUN, J.; VEGA, F. E. Fungal endophytes in germinated seeds of the common bean, *Phaseolus vulgaris*. **Fun. Bio.**, v. 120, p. 783-790, 2016.

STANGARLIN, J. R; SCHWAN-ESTRADA, K. R. F; SILVA CRUZ, M. E; NOZAKI, M. H. Plantas Mediciniais e Controle Alternativo de Fitopatógenos. **Biotecnol. Ciênc. Desenv.**, Brasília, n.11, p.16-21, 1999.

THANGAVELU, R; SUNDARARAJU, P; SATHIAMOORTHY, S. Management of Anthracnose Disease of Banana Caused by *Colletotrichum Musae* Using Plant Extracts. **J. Hort. Sci. Biotechnol.**, Dundee, v. 79, p. 664–668, 2004.

VALADARES, R. B. S; PEREIRA, M. C; KASUYA, M. C. M; CARDOSO, E. J. B. N. Isolamento e Identificação de Fungos Micorrízicos de *Gomezia sp.* (Orchidaceae) em uma Floresta de Araucária do Estado de São Paulo. In: Fertbio, 2008, Londrina. **Anais...**, p. 255-258, 2008.

VOGT, V; CIFUENTE, D; TONN, C; SABINI, L; ROSAS, S. Antifungal Activity *In Vitro* and *In vivo* of Extracts and Lignans Isolated from *Larrea Divaricata* Cav. Against Phytopathogenic Fungus. **Ind. Crops Prod.**, St Martin d'Herès, v. 42, p. 583– 586, 2012.